



Williams College ECON 523:

Program Evaluation for International Development

Lecture 5: Two-Way Fixed Effects

Professor: Pamela Jakiela

photo: Daniella Van Leggelo Padilla / World Bank

Variation in Treatment Timing

Example: States Adopted Medicaid at Different Times




Figure 2.
Medicaid Adoption by Quarter
Notes: Adoption dates come from the Department of Health Education and Welfare (1970) & Social Security Administration (2013). The map is shaded relative to the quarter of adoption and states are labeled with the month and year of adoption.

source: Boudreaux, Golberstein, and McAlpine (*Journal of Health Economics*, 2016)

Example: Counties Opening Community Health Centers




FIGURE 3. ESTABLISHMENT OF COMMUNITY HEALTH CENTERS BY COUNTY OF SERVICE DELIVERY, 1965–1980

Note: Dates are the first year that a CHC was established in the county.

Source: Information on CHCs drawn from NACAP and PHS reports.

source: Bailey and Goodman-Bacon (*AER*, 2015)

Example: African Countries Introducing Multi-Party Elections



FIGURE A.2. Geographical distribution of democratized countries since 1990. Black-colored countries are democratized since 1990; grey-colored countries are the other countries in the sample for infant mortality analysis. Democratized countries include the Comoros, tiny islands to the northwest of Madagascar, which may not be visible as black-colored.

source: Kudamatsu (JEEA, 2012)

Two-Way Fixed Effects Estimates of β^{DD}

What exactly is β^{DD} ?

$$Y_{it} = \alpha_i + \gamma_t + \beta^{DD} D_{it} + \varepsilon_{it}$$

unit fixed effects

time fixed effects

treatment dummy
that turns "on" at
different times


What exactly is β^{DD} in TWFE?

	$t = 1$	$t = 2$	$t = 3$	$t = 4$	$t = 5$	
ID 1	0	0	1	1	1	treatment
ID 2	0	0	1	1	1	
ID 3	0	0	0	0	0	comparison
ID 4	0	0	0	0	0	

Multiple Treatment and Comparison Groups


	$t = 1$	$t = 2$	$t = 3$	$t = 4$	$t = 5$	$t = 6$	$t = 7$	$t = 8$
ID 1	0	0	1	1	1	1	1	1
ID 2	0	0	1	1	1	1	1	1
ID 3	0	0	0	0	0	1	1	1
ID 4	0	0	0	0	0	1	1	1
ID 5	0	0	0	0	0	0	0	0
ID 6	0	0	0	0	0	0	0	0

Decomposition into Timing Groups




Panel with variation in treatment timing can be decomposed into distinct **timing groups** reflecting observed start of treatment

Decomposition into Timing Groups



Example: with three timing groups (one of which is never treated), can construct three timing windows (pre, middle, post or $t = 1, 2, 3$)


Decomposition into Standard 2×2 DDs



Economics 523 (Professor Jakiela)

Two-Way Fixed Effects, Slide 15

Decomposition into Standard 2×2 DDs




We know DD estimate of treatment effect for each timing group:

$$\begin{aligned}\hat{\beta}_{AC}^{DD} &= (\bar{Y}_A^{POST} - \bar{Y}_C^{POST}) - (\bar{Y}_A^{PRE} - \bar{Y}_C^{PRE}) \\ &= (\bar{Y}_A^{t=2,3} - \bar{Y}_C^{t=2,3}) - (\bar{Y}_A^{t=1} - \bar{Y}_C^{t=1})\end{aligned}$$

Economics 523 (Professor Jakiela)

Two-Way Fixed Effects, Slide 16

Forbidden Comparisons



Some 2×2 diff-in-diffs (implicitly) use already-treated groups as the comparison

Two-Way Fixed Effects β^{DD} as a Weighted Sum

The two-way fixed effects estimator β^{DD} is a weighted sum of 2×2 diff-in-diff estimators across all possible pairwise combinations of timing groups (Goodman-Bacon 2021)

- Some use an already-treated group as comparison
 - ▶ Creates problems if treatment effect grows/changes over time
 - ▶ TWFE imposes a model of homogeneous treatment effects
 - ▶ When treatment effects evolve over time, model is mis-specified

We can use Frisch-Waugh-Lovell to construct the TWFE/OLS weights used to generate β^{DD}

- Weights on treated units are not always positive (they are also used as comparison)

Two-Way Fixed Effects as Univariate Regression

Two-way fixed effects is equivalent to univariate regression:

$$\tilde{Y}_{it} = \alpha + \tilde{D}_{it} + \epsilon_{it}$$

where $\tilde{Y}_{it} = Y_{it} - \bar{Y}_t - (\bar{Y}_i - \bar{\bar{Y}})$ and \tilde{D}_{it} defined analogously

↑
“grand mean”

(just the mean across i and t)

Two-Way Fixed Effects as Univariate Regression

Two-way fixed effects is equivalent to univariate regression:

$$\tilde{Y}_{it} = \alpha + \tilde{D}_{it} + \epsilon_{it}$$

where $\tilde{Y}_{it} = Y_{it} - \bar{Y}_t - (\bar{Y}_i - \bar{\bar{Y}})$ and \tilde{D}_{it} defined analogously

⇒ Treatment dummy now effectively continuous measure \tilde{D}_{it}

$$\hat{\beta}^{OLS} = \sum_{it} \tilde{Y}_{it} \underbrace{\left(\frac{\tilde{D}_{it} - \bar{\tilde{D}}_{it}}{\sum_i (\tilde{D}_{it} - \bar{\tilde{D}}_{it})^2} \right)}_{\text{OLS weight}}$$

Two-Way Fixed Effects as Univariate Regression

Two-way fixed effects is equivalent to univariate regression:

$$\tilde{Y}_{it} = \alpha + \tilde{D}_{it} + \epsilon_{it}$$

where $\tilde{Y}_{it} = Y_{it} - \bar{Y}_t - (\bar{Y}_i - \bar{\bar{Y}})$ and \tilde{D}_{it} defined analogously

⇒ Treatment dummy now effectively continuous measure \tilde{D}_{it}

$$\hat{\beta}^{OLS} = \sum_{it} \tilde{Y}_{it} \underbrace{\left(\frac{\tilde{D}_{it} - \bar{\tilde{D}}_{it}}{\sum_i (\tilde{D}_{it} - \bar{\tilde{D}}_{it})^2} \right)}_{\text{OLS weight}} \text{ where } \bar{\tilde{D}}_{it} = 0$$

Diff-in-Diff without Staggered Treatment Timing: Review

	$t = 1$	$t = 2$	$t = 3$	$t = 4$
ID 1	0	1	1	1
ID 2	0	0	0	0

Diff-in-Diff without Staggered Treatment Timing: Review

$$D_{it} - \bar{D}_t - (\bar{D}_i - \bar{\bar{D}})$$

	$t = 1$	$t = 2$	$t = 3$	$t = 4$	
ID 1	-0.375	0.125	0.125	0.125	← Equal weight
ID 2	0.375	-0.125	-0.125	-0.125	

Diff-in-Diff with Staggered Treatment Timing: Example

	$t = 1$	$t = 2$	$t = 3$	$t = 4$
ID 1	0	1	1	1
ID 2	0	0	0	1
\bar{D}_t	0	0.5	0.5	1

mean treatment in period t

Diff-in-Diff with Staggered Treatment Timing: Example

	Y_{it}			
	$t = 1$	$t = 2$	$t = 3$	$t = 4$
ID 1	0	10	10	10
ID 2	0	0	0	10

Let $Y_{it} = \gamma_i + \lambda_t + \delta_{it}$

Treated cells

Positive weights (in treatment group)

Diff-in-Diff with Staggered Treatment Timing: Example

	Y_{it}			
	$t = 1$	$t = 2$	$t = 3$	$t = 4$
ID 1	0	10	10	10
ID 2	0	0	0	10

$\hat{\beta}_{OLS} = 10$

homogeneous impacts:
 $E[\hat{\beta}_{OLS}] = ATE$

Treated cells

Positive weights (in treatment group)


Diff-in-Diff with Staggered Treatment Timing: Example

	Y_{it}				$\hat{\beta}_{OLS} = 6$
	$t = 1$	$t = 2$	$t = 3$	$t = 4$?
ID 1	0	2	2	2	
ID 2	0	0	0	10	

Treated cells

Positive weights (in treatment group)

Diff-in-Diff with Staggered Treatment Timing: Example


	Y_{it}				$\hat{\beta}_{OLS} = -2$
	$t = 1$	$t = 2$	$t = 3$	$t = 4$	
ID 1	0	2	2	10	
ID 2	0	0	0	2	

Treated cells

Positive weights (in treatment group)

TWFE in Practice

Policy Context: Free Primary Education in Sub-Saharan Africa



TWFE Specification: Countries that Implemented Free Primary

$$\text{TWFE specification: } Primary_{it} = \alpha_i + \gamma_t + \beta FPE_{it} + \varepsilon_{it}$$

Stata code: `reg primary fpe i.cid i.year, cluster(cid)`



	(1)
	Enrollment
Free primary education	20.428 (9.120) [0.042]
Country fixed effects	Yes
Year fixed effects	Yes
Never treated	No

Dependent variable: gross enrollment ratio.
Data on gross enrollment ratio in 15 countries comes from the World Development Indicators, years 1981 through 2015. Standard errors (clustered at the country level) in parentheses; p-values in square brackets.


TWFE Diagnostics

1. Are treated observations getting negative weight in my TWFE estimation?
 - ▶ Are treated observations (i.e. country-years) being weighted in a sensible way?
2. Are treatment effects (likely to be) heterogeneous? If yes, how?
 - ▶ Conceptually: do you expect the treatment effects to vary over time, across units, or both?
 - ▶ Do you see evidence contradicting the assumption of homogeneous treatment effects?
 - ▶ Event study specifications
 - ▶ Scatter plots of residuals
 - ▶ Are your estimated treatment effects robust across specifications?

Negative Weights





Including Never Treated Countries to Eliminate Negative Weights







	(1)	(2)
	Enrollment	Enrollment
Free primary education	20.428 (9.120) [0.042]	18.951 (6.733) [0.008]
Country fixed effects	Yes	Yes
Year fixed effects	Yes	Yes
Never treated	No	Yes

Dependent variable: gross enrollment ratio. Data on gross enrollment ratio in 15 countries comes from the World Development Indicators, years 1981 through 2015. Standard errors (clustered at the country level) in parentheses; p-values in square brackets.

Including Never Treated Countries to Eliminate Negative Weights



Including Never Treated Countries to Eliminate Negative Weights



Event Study Specifications

Negative weights are a major issue if treatment effects change over (relative) time

- Relative time is the number of years since treatment was implemented (in country t)
- We can also think of negative relative time as years until treatment starts (in country t)

An **event study** specification allows us to estimate treatment effects for every (relative) time

- Provides direct evidence on the stability of the treatment effect (over time)
- Also allows us to check for violations of common (pre)trends
- Because we are estimating many parameters instead of one, statistical power is an issue

Event Study Specifications

Let G_i indicate the time t when treatment starts in country i



$\Rightarrow R_{it} = t - G_i$ is relative time, and treatment starts when $R_{it} = 0$

TWFE event study specification:



$$Primary_{it} = \alpha_i + \gamma_t + \sum_{r \leq -2} \beta_r 1[R_{it} = r] + \sum_{r \geq 0} \delta_r 1[R_{it} = r] + \varepsilon_{it}$$

Impacts are defined relative to $R_{it} = -1$, the last period before treatment


The Impacts of Free Primary: Event Study Specifications



Plotting the Residuals



Trimming the Sample



Other Robustness Checks

Summarizing these and other approaches to assessing the robustness of TWFE estimates:

- Trim sample to eliminate negative weights and common trends violations, focus on post-treatment window when assumption of constant treatment effects seems justifiable
- Jackknife estimation: drop one country at a time to assess other types of heterogeneity
- Include unit-specific trends as controls
- Other Stata packages: `csdid`, `eventstudyinteract` (see Roth et al. 2023 for more)

With homogeneous treatment effects, these approaches should generate similar results

TWFE: Checklist

- Check for negative weights, and consider eliminating them
 - ▶ The most important thing is to know what you are estimating
- Assess the linearity of the residuals: is homogeneity a reasonable assumption?
- Implement an event study design, if feasible given sample/power
- Robustness checks, more robustness checks, and even more robustness checks